arXiv:math/0107134v2 [math.AG] 10 Aug 2001

MOTIVIC INTEGRATION ON SMOOTH RIGID VARIETIES AND INVARIANTS OF DEGENERATIONS

by

François Loeser & Julien Sebag

1. Introduction

In the last years, motivic integration has shown to be a quite powerful tool in producing new invariants in birational geometry of algebraic varieties over a field k, say of characteristic zero, cf. [**17**][**2**][**3**][**10**][**11**][**12**]. Let us explain the basic idea behind such results. If $h : Y \to X$ is a proper birational morphism between k-algebraic varieties, the induced morphism $\mathcal{L}(Y) \to \mathcal{L}(X)$ between arc spaces (cf. [**10**]) is an isomorphism outside subsets of infinite codimension. By a fundamental change of variable formula, motivic integrals on $\mathcal{L}(X)$ may be computed on $\mathcal{L}(Y)$ when Y is smooth.

In the present paper we develop similar ideas in the somewhat dual situation of degenerating families over complete discrete valuation rings with perfect residue field, for which rigid geometry appears to be a natural framework. More precisely, let R be a complete discrete valuation ring with fraction field K and perfect residue field k. We construct a theory of motivic integration for smooth$^{(1)}$ rigid K-spaces, always assumed to be quasi-compact and separated. Let X be a smooth rigid K-space of dimension d. Our construction assigns to a gauge form ω on X, *i.e.* a nowhere vanishing differential form of degree d on X, an integral $\int_X \omega d\tilde{\mu}$ with value in the ring $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$. Here $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$ is the localization with respect to the class of the affine line of the Grothendieck group of algebraic varieties over k. In concrete terms, two varieties over k define the same class in $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$ if they become isomorphic after cutting them into locally closed pieces and stabilization by product with a power of the affine line. More generally, if ω is a differential form of degree d on X, we define an integral $\int_X \omega d\mu$ with value in the ring $\widehat{K_0(\mathrm{Var}_k)}$, which is the completion of $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$ with respect to the filtration by virtual dimension (see § 3.4). The construction is done by viewing X as the generic fibre of some formal R-scheme $\mathcal{X}$. To such a formal R-scheme, by mean of the Greenberg functor $\mathcal{X} \mapsto \mathrm{Gr}(\mathcal{X})$ one associates a certain k-scheme $\mathrm{Gr}(\mathcal{X})$, which, when $R = k[[t]]$, and $\mathcal{X}$ is the formal completion of $X_0 \otimes k[[t]]$, for X_0 an algebraic variety over k, is nothing else that the arc space $\mathcal{L}(X_0)$ considered before. We may then use the general theory of motivic integration on schemes $\mathrm{Gr}(\mathcal{X})$ which is developed in [**24**]. Of course, for the construction to work one needs to check that it is independent of the chosen model. This is done by using two main ingredients: the theory of weak Néron models developed in [**5**] and [**9**], and the analogue for schemes of the form $\mathrm{Gr}(\mathcal{X})$ of the change of variable formula which is proven in [**24**]. In fact the theory of weak Néron models really pervades the whole paper and some parts of the book [**5**] were crying out for their use in motivic integration(2).

---

(1) The extension to singular rigid spaces when K is of characteristic zero will be considered in a separate publication.

As an application of our theory, we are able to assign in a canonical way to any smooth quasi-compact and separated rigid K-space X an element $\lambda(X)$ in the quotient ring $K_0(\mathrm{Var}_k)_{\mathrm{loc}}/(\mathbf{L}-1)K_0(\mathrm{Var}_k)_{\mathrm{loc}}$, where $\mathbf{L}$ stands for the class of the affine line. When X admits a formal R-model with good reduction, $\lambda(X)$ is just the class of the fibre of that model. More generally, if $\mathcal{U}$ is a weak Néron model of X, $\lambda(X)$ is equal to the class of the special fibre of $\mathcal{U}$ in $K_0(\mathrm{Var}_k)_{\mathrm{loc}}/(\mathbf{L}-1)K_0(\mathrm{Var}_k)_{\mathrm{loc}}$. In particular it follows that this class is independent of the choice of the weak Néron model $\mathcal{U}$.

This result can be viewed as a rigid analogue of a result of Serre concerning compact smooth locally analytic varieties over a local field. To such a variety $\tilde{X}$, Serre associates in [**25**], using classical p-adic integration, an invariant $s(\tilde{X})$ in the ring $\mathbf{Z}/(q-1)\mathbf{Z}$, where q denotes the cardinality of the finite field k. Counting rational points in k yields a canonical morphism $K_0(\mathrm{Var}_k)_{\mathrm{loc}}/(\mathbf{L}-1)K_0(\mathrm{Var}_k)_{\mathrm{loc}} \to \mathbf{Z}/(q-1)\mathbf{Z}$, and we show that the image by this morphism of our motivic invariant $\lambda(X)$ of a smooth rigid K-space X is equal to the Serre invariant of the underlying locally analytic variety.

Unless making additional assumptions on X, one cannot hope to lift our invariant $\lambda(X)$ to a class in the Grothendieck ring $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$, which would be a substitute for the class of the special fibre of **the** Néron model, when such a Néron model happens to exist. In the particular situation where X is the analytification of a Calabi-Yau variety over K, *i.e.* a smooth projective algebraic variety over K of pure dimension d with Ω^d_X trivial, this can be achieved: one can attach to X a canonical element of $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$, which, if X admits a proper and smooth R-model $\mathcal{X}$, is equal to the class of the special fibre $\mathcal{X}_0$ in $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$. In particular, if X admits two such models $\mathcal{X}$ and $\mathcal{X}'$, the class of the special fibres $\mathcal{X}_0$ and $\mathcal{X}'_0$ in $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$ are equal, which may be seen as an analogue of Batyrev's result on birational Calabi-Yau varieties [**1**].

The paper is organized as follows. Section 2 is devoted to preliminaries on formal schemes, the Greenberg functor and weak Néron models. In the following section, we review the results on motivic integration on formal schemes obtained by the second named author in [**24**] which are needed in the present work. We are then able in section 4 to construct a motivic integration on smooth rigid varieties and to prove the main results which were mentionned in the present introduction. Finally, in section 5, guided by the analogy with arc spaces, we formulate an analogue of the Nash problem, which is about the relation between essential (*i.e.* appearing in every resolution) components of resolutions of a singular variety and irreducible components of spaces of truncated arcs on the variety, for formal R-schemes with smooth generic fibre. In this context, analogy suggests there might be some relation between essential components of weak Néron models of a given formal R-scheme $\mathcal{X}$ with smooth generic fibre and irreducible components of the truncation $\pi_n(\mathrm{Gr}(\mathcal{X}))$ of its Greenberg space for $n \gg 0$. As a very first step in that direction we compute the dimension of the contribution of a given irreducible component to the truncation.

---

(2) cf. the remark at the bottom of p.105 in [**5**].

2. Preliminaries on formal schemes and Greenberg functor

2.1. Formal schemes. — In this paper R will denote a complete discrete valuation ring with residue field k and fraction field K. We shall assume k is perfect. We shall fix once for all an uniformizing parameter ϖ and we shall set $R_n := R/(\varpi)^{n+1}$, for $n \geq 0$. In the whole paper, by a formal R-scheme, we shall always mean a quasi-compact, separated, locally topologically of finite type formal R-scheme, in the sense of § 10 of [**13**]. A formal R-scheme is a locally ringed space $(\mathcal{X}, \mathcal{O}_{\mathcal{X}})$ in topological R-algebras. It is equivalent to the data, for every $n \geq 0$, of the R_n-scheme $X_n = (\mathcal{X}, \mathcal{O}_{\mathcal{X}} \otimes_R R_n)$. The k-scheme X_0 is called the special fibre of $\mathcal{X}$. As a topological space $\mathcal{X}$ is isomorphic to X_0 and $\mathcal{O}_{\mathcal{X}} = \varprojlim \mathcal{O}_{X_n}$. We have $X_n = X_{n+1} \otimes_{R_{n+1}} R_n$ and $\mathcal{X}$ is canonically isomorphic to the inductive limit of the schemes X_n in the category of formal schemes. Locally $\mathcal{X}$ is isomorphic to an affine formal R-scheme of the form $\mathrm{Spf} A$ with A an R-algebra topologically of finite type, *i.e.* a quotient of a restricted formal series algebra $R\{T_1, \ldots, T_m\}$. If $\mathcal{Y}$ and $\mathcal{X}$ are R-formal schemes, we denote by $\mathrm{Hom}_R(\mathcal{Y}, \mathcal{X})$ the set of morphisms of formal R-schemes $\mathcal{Y} \to \mathcal{X}$, *i.e.* morphisms between the underlying locally topologically ringed spaces over R (cf. § 10 of [**13**]). It follows from Proposition 10.6.9 of [**13**], that the canonical morphism $\mathrm{Hom}_R(\mathcal{Y}, \mathcal{X}) \to \varprojlim \mathrm{Hom}_{R_n}(\mathcal{Y}_n, \mathcal{X}_n)$ is a bijection.
If k is a field, by a variety over k we mean a separated reduced scheme of finite type over k.

2.2. Extensions. — Let A be a k-algebra. We set $L(A) = A$ when R is a ring of equal characteristic and $L(A) = W(A)$, the ring of Witt vectors, when R is a ring of unequal characteristic and we denote by R_A the ring $R_A := R \otimes_{L(k)} L(A)$. When F is a field containing k, we denote by K_F the field of fractions of R_F. When the field F is perfect, the ring R_F is a discrete valuation ring and, furthermore, the uniformizing parameter ϖ in R induces an uniformizing parameter in R_F. Hence, since k is assumed to be perfect, the extension $R \to R_F$ has ramification index 1 in the terminology of § 3.6 of [**5**].

2.3. The Greenberg Functor. — We shall recall some material from [**14**] and § 9.6 of [**5**]. Let us remark, that, when R is a ring of equal characteristic, we can view R_n as the set of k-valued points of some affine space $\mathbf{A}_k^m$ which we shall denote by $\mathcal{R}_n$, in a way compatible with the k-algebra structure. When R is a ring of unequal characteristic, R_n can no longer be viewed as a k-algebra. However, using Witt vectors, we may still interpret R_n as the set of k-valued points of a ring scheme $\mathcal{R}_n$, which, as a k-scheme, is isomorphic to some affine space $\mathbf{A}_k^m$. Remark we have canonical morphisms $\mathcal{R}_{n+1} \to \mathcal{R}_n$.

Now, for every $n \geq 0$, we consider the functor h_n^* which to a k-scheme T associates the locally ringed space $h_n^*(T)$ which has T as underlying topological space and $\mathcal{H}om_k(T, \mathcal{R}_n)$ as structure sheaf. In particular, for any k-algebra A,

$$h_n^*(A) = \mathrm{Spec}(R_n \otimes_{L(k)} L(A)).$$

Taking $A = k$, we see that $h_n^* T$ is a locally ringed space over $\mathrm{Spec}\, R_n$.

By a fundamental result of Greenberg [**14**] (which in the equal characteristic case amounts to Weil restriction of scalars), for R_n-schemes X_n, locally of finite type, the functor

$$T \longmapsto \mathrm{Hom}_{R_n}(h_n^*(T), X_n)$$

from the category of k-schemes to the category of sets is represented by a k-scheme $\mathrm{Gr}_n(X_n)$ which is locally of finite type. Hence, for any k-algebra A,

$$\mathrm{Gr}_n(X_n)(A) = X_n(R_n \otimes_{L(k)} L(A)),$$

and, in particular, setting $A = k$, we have

$$\mathrm{Gr}_n(X_n)(k) = X_n(R_n).$$

Among basic properties the Greenberg functor respects closed immersions, open imersions, fibred products, smooth and étale morphisms and also sends affines to affines.

Now let us consider again a formal R-scheme $\mathcal{X}$. The canonical adjunction morphism $h_{n+1}^*(\mathrm{Gr}_{n+1}(X_{n+1})) \to X_{n+1}$ gives rise, by tensoring with R_n, to a canonical morphism of of R_n-schemes $h_n^*(\mathrm{Gr}_{n+1}(X_{n+1})) \to X_n$, from which one derives, again by adjunction, a canonical morphism of k-schemes

$$\theta_n : \mathrm{Gr}_{n+1}(X_{n+1}) \longrightarrow \mathrm{Gr}_n(X_n).$$

In this way we attach to the formal scheme $\mathcal{X}$ a projective system $(\mathrm{Gr}_n(X_n))_{n \in \mathbf{N}}$ of k-schemes. The transition morphisms θ_n being affine, the projective limit

$$\mathrm{Gr}(\mathcal{X}) := \varprojlim \mathrm{Gr}_n(X_n)$$

exists in the category of k-schemes.

Let T be a k-scheme. We denote by $h^*(T)$ the locally ringed space which has T as underlying topological space and $\varprojlim \mathcal{H}om_k(T, \mathcal{R}_n)$ as structure sheaf. It is a locally ringed space over $\mathrm{Spf}\, R$ which identifies with the projective limit of the spaces $h_n^*(T)$ in the category of locally ringed spaces. Furthermore one checks, similarly

as in Proposition 10.6.9 of [**13**], that the canonical morphism $\mathrm{Hom}_R(h^*(T), \mathcal{X}) \to \varprojlim \mathrm{Hom}_{R_n}(h_n^*(T), \mathcal{X}_n)$ is a bijection for every formal R-scheme $\mathcal{X}$.
Putting everything together we get the following:

2.3.1. Proposition. — *Let $\mathcal{X}$ be a quasi-compact, locally topologically of finite type formal R-scheme. The functor*

$$T \longmapsto \mathrm{Hom}_R(h^*(T), \mathcal{X})$$

from the category of k-schemes to the category of sets is represented by the k-scheme $\mathrm{Gr}(\mathcal{X})$. □

In particular, for every field F containing k, there are canonical bijections

$$\mathrm{Gr}(\mathcal{X})(F) \simeq \mathrm{Hom}_R(\mathrm{Spf}\, R_F, \mathcal{X}) \simeq \mathcal{X}(R_F).$$

One should note that, in general, $\mathrm{Gr}(\mathcal{X})$ is not of finite type, even if $\mathcal{X}$ is a quasi-compact, topologically of finite type formal R-scheme.
In this paper, we shall always consider the schemes $\mathrm{Gr}_n(X_n)$ and $\mathrm{Gr}(\mathcal{X})$ with their reduced structure.
Sometimes, by abuse of notation, we shall write $\mathrm{Gr}_n(\mathcal{X})$ for $\mathrm{Gr}_n(X_n)$.

2.3.2. Proposition. —

1. *The functor* Gr *respects open and closed immersions, fibre products, and sends affine formal R-schemes to affine k-schemes.*
2. *Let $\mathcal{X}$ be a formal quasi-compact and separated R-scheme and let $(\mathcal{O}_i)_{i \in J}$ be a finite covering by formal open subschemes. There are canonical isomorphisms* $\mathrm{Gr}(\mathcal{O}_i \cap \mathcal{O}_j) \simeq \mathrm{Gr}(\mathcal{O}_i) \cap \mathrm{Gr}(\mathcal{O}_j)$ *and the scheme* $\mathrm{Gr}(\mathcal{X})$ *is canonically isomorphic to the scheme obtained by glueing the schemes* $\mathrm{Gr}(\mathcal{O}_i)$.

Proof. — Assertion (1) for the functor Gr_n is proved in [**14**] and [**5**], and follows for Gr by taking projective limits. Assertion (2) follows from (1) and the universal property defining Gr. □

2.3.3. Remark. — Assume we are in the equal characteristic case, *i.e.* $R = k[[\varpi]]$. For X an algebraic variety over k, we can consider the formal R-scheme $X \widehat{\otimes} R$ obtained by base change and completion. We have canonical isomorphisms $\mathrm{Gr}(X \widehat{\otimes} R) \simeq \mathcal{L}(X)$ and $\mathrm{Gr}_n(X \otimes R_n) \simeq \mathcal{L}_n(X)$, where $\mathcal{L}(X)$ and $\mathcal{L}_n(X)$ are the arc spaces considered in [**10**].

2.4. Smoothness. — Let us recall the definition of smoothness for morphisms of formal R-schemes. A morphism $f : \mathcal{X} \to \mathcal{Y}$ of formal R-schemes is smooth at a point x of X_0 of relative dimension d if it is flat at x and the induced morphism $f_0 : X_0 \to Y_0$ is smooth at x of relative dimension d. An equivalent condition (cf. Lemma 1.2 of [**7**]) is that for every n in $\mathbf{N}$ the induced morphism $f_n : X_n \to Y_n$ is smooth at x of relative dimension d. The morphism f is smooth if it is smooth at every point of X_0. The formal R-scheme $\mathcal{X}$ is smooth at a point x of X_0 if the structural morphism is smooth at x.

Let $\mathcal{X}$ be a flat formal R-scheme of relative dimension d. We denote by $\mathcal{X}_{\mathrm{sing}}$ the closed formal subscheme of $\mathcal{X}$ defined by the radical of the Fitting ideal sheaf $\mathrm{Fitt}_d \Omega_{\mathcal{X}/R}$. The formal R-scheme $\mathcal{X}$ is smooth at a point x of X_0 (resp. is smooth) if and only if x is not in $\mathcal{X}_{\mathrm{sing}}$ (resp. $\mathcal{X}_{\mathrm{sing}}$ is empty).

2.5. Greenberg's Theorem. — The following statement, which is an adaptation of a result of Schappacher [**23**], is an analogue of Greenberg's Theorem [**15**] in the framework of formal schemes. We refer to [**24**] for a more detailled exposition.

2.5.1. ***Theorem.*** — *Let R be a complete discrete valuation ring and let $\mathcal{X}$ be a formal R-scheme. For every $n \geq 0$, there exists an integer $\gamma_{\mathcal{X}}(n) \geq n$ such that, for every field F containing k, and every x in $\mathcal{X}(R_F/\varpi^{\gamma_{\mathcal{X}}(n)+1})$, the image of x in $\mathcal{X}(R_F/\varpi^{n+1})$ may be lifted to a point in $\mathcal{X}(R_F)$.*

2.5.2. ***Remark.*** — The function $n \mapsto \gamma_{\mathcal{X}}(n)$ is called the Greenberg function of $\mathcal{X}$.

2.6. Rigid spaces. — For $\mathcal{X}$ a flat formal R-scheme we shall denote by $\mathcal{X}_K$ its generic fibre in the sense of Raynaud [**22**]. By Raynaud's Theorem [**22**], [**6**], the functor $\mathcal{X} \mapsto \mathcal{X}_K$ induces an equivalence between the localization of the category of quasi-compact flat formal R-schemes by admissible formal blowing-ups, and the category of rigid K-spaces which are quasi-compact and quasi-separated. Furthemore, $\mathcal{X}$ is separated if and only if $\mathcal{X}_K$ is separated (cf. Proposition 4.7 of [**6**]). Recall that for the blowing-up of an ideal sheaf $\mathcal{I}$ to be admissible means that $\mathcal{I}$ contains some power of the uniformizing parameter ϖ. In the paper all rigid K-spaces will be assumed to be quasi-compact and separated.

2.7. Weak Néron models. — We shall denote by R^{sh} a strict henselization of R and by K^{sh} its field of fractions.

2.7.1. ***Definition.*** — Let X be a smooth rigid K-variety. A weak formal Néron model(3) of X is a smooth formal R-scheme $\mathcal{U}$, whose generic fibre $\mathcal{U}_K$ is an open rigid subspace of X_K, and which has the property that the canonical map $\mathcal{U}(R^{\mathrm{sh}}) \to X(K^{\mathrm{sh}})$ is bijective.

The construction of weak Néron models using Néron's smoothening process presented in [**5**], carries over almost literally from R-schemes to formal R-schemes, and gives, as explained in [**9**], the following result:

2.7.2. ***Theorem.*** — *Let $\mathcal{X}$ be a quasi-compact formal R-scheme, whose generic fibre $\mathcal{X}_K$ is smooth. Then there exists a morphism of formal R-schemes $\mathcal{X}' \to \mathcal{X}$, which is the composition of a sequence of formal blowing-ups with centers in the corresponding special fibres, such that every R^{sh}-valued point of $\mathcal{X}$ factors through the smooth locus of $\mathcal{X}'$.*

---

(3) We follow here the terminology of [**9**] which is somewhat different from that of [**5**].

One deduces the following omnibus statement:

2.7.3. Proposition. — *Let X be a smooth quasi-compact and separated rigid K-space and let $\mathcal{X}$ be a formal R-model of X, i.e a quasi-compact formal R-scheme $\mathcal{X}$ with generic fibre X. Then there exists a weak formal Néron model $\mathcal{U}$ of X which dominates $\mathcal{X}$ and which is quasi-compact. Furthermore the canonical map $\mathcal{U}(R^{\mathrm{sh}}) \to \mathcal{X}(R^{\mathrm{sh}})$ is a bijection and for every perfect field F containing k, the formal R_F-scheme $\mathcal{U} \otimes_R R_F$ is a weak Néron model of the rigid K_F-space $X \otimes_K K_F$. In particular, the morphism $\mathcal{U} \to \mathcal{X}$ induces a bijection between points of $\mathrm{Gr}(\mathcal{U})$ and $\mathrm{Gr}(\mathcal{X})$.*

Proof. — We choose a formal model $\mathcal{X}$ of X such that we are in the situation of Theorem 2.7.2. The smooth locus $\mathcal{U}$ of $\mathcal{X}'$ is quasi-compact and is a weak Néron model of $\mathcal{X}_K$, since, by [**9**] 2.2 (ii), every K^{sh}-valued point of $\mathcal{X}_K$ extends uniquely to a R^{sh}-valued point of $\mathcal{X}$. Also, it follows from Corollary 6 of § 3.6 [**5**] that, if $\mathcal{U}$ is a weak Néron model of the rigid K-space X, then for every field F containing k, the formal R_F-scheme $\mathcal{U} \otimes_R R_F$ is a weak Néron model of the rigid K_F-space $X \otimes_K K_F$. □

3. Motivic integration on formal schemes

The material in this section is borrowed from [**24**] to which we shall refer for details.

3.1. Truncation. — For $\mathcal{X}$ a formal R-scheme, we shall denote by $\pi_{n,\mathcal{X}}$ or π_n the canonical projection $\mathrm{Gr}(\mathcal{X}) \to \mathrm{Gr}_n(X_n)$, for n in $\mathbf{N}$.
Let us first state the following corollary of Theorem 2.5.1:

3.1.1. Proposition. — *Let $\mathcal{X}$ be a formal R-scheme. The image $\pi_n(\mathrm{Gr}(\mathcal{X}))$ of $\mathrm{Gr}(\mathcal{X})$ in $\mathrm{Gr}_n(X_n)$ is a constructible subset of $\mathrm{Gr}_n(X_n)$. More generally, if C is a constructible subset of $\mathrm{Gr}_m(X_m)$, $\pi_n(\pi_m^{-1}(C))$ is a constructible subset of $\mathrm{Gr}_n(X_n)$, for every $n \geq 0$.*

Proof. — Indeed, it follows from Theorem 2.5.1 that $\pi_n(\mathrm{Gr}(\mathcal{X}))$ is equal to the image of $\mathrm{Gr}_{\gamma(n)}(X_{\gamma(n)})$ in $\mathrm{Gr}_n(X_n)$. The morphism $\mathrm{Gr}_{\gamma(n)}(X_{\gamma(n)}) \to \mathrm{Gr}_n(X_n)$ being of finite type, first the statement follows from Chevalley's Theorem. For the second statement, one may assume $m = n$, and the proof proceeds as before. □

3.1.2. Proposition. — *Let $\mathcal{X}$ be a smooth formal separated R-scheme (quasi-compact, locally topologically of finite type over R), of relative dimension d.*

1. *For every n, the morphism $\pi_n : \mathrm{Gr}(\mathcal{X}) \to \mathrm{Gr}_n(X_n)$ is surjective.*
2. *For every n and m in $\mathbf{N}$, the canonical projection $\mathrm{Gr}_{n+m}(X_{n+m}) \to \mathrm{Gr}_n(X_n)$ is a locally trivial fibration for the Zariski topology with fibre $\mathbf{A}_k^{dm}$.*

We say that a map $\pi : A \to B$ is a piecewise morphism if there exists a finite partition of the domain of π into locally closed subvarieties of X such that the restriction of π to any of these subvarieties is a morphism of schemes.

3.2. Away from the singular locus. —

Let $\mathcal{X}$ be a formal R-scheme and consider its singular locus $\mathcal{X}_{\mathrm{sing}}$ defined in 2.4. For any integer $e \geq 0$, we view $\mathrm{Gr}_e(\mathcal{X}_{\mathrm{sing},e})$ as contained in $\mathrm{Gr}_e(\mathcal{X})$ and we set

$$\mathrm{Gr}^{(e)}(\mathcal{X}) := \mathrm{Gr}(\mathcal{X}) \setminus \pi_e^{-1}(\mathrm{Gr}_e(\mathcal{X}_{\mathrm{sing},e})).$$

We say that a map $\pi : A \to B$ between k-constructible sets A and B is a piecewise trivial fibration with fibre F, if there exists a finite partition of B in subsets S which are locally closed in Y such that $\pi^{-1}(S)$ is locally closed in X and isomorphic, as a variety over k, to $S \times F$, with π corresponding under the isomorphism to the projection $S \times F \to S$. We say that the map π is a piecewise trivial fibration over some constructible subset C of B, if the restriction of π to $\pi^{-1}(C)$ is a piecewise trivial fibration onto C.

3.2.1. Proposition. — *Let $\mathcal{X}$ a flat formal R-scheme of relative dimension d. There exists an integer $c \geq 1$ such that, for every integers e and n in $\mathbf{N}$ such that $n \geq ce$, the projection*

$$\pi_{n+1}(\mathrm{Gr}(\mathcal{X})) \longrightarrow \pi_n(\mathrm{Gr}(\mathcal{X}))$$

is a piecewise trivial fibration over $\pi_n(\mathrm{Gr}^{(e)}(\mathcal{X}))$ with fibre $\mathbf{A}_k^d$.

3.3. Dimension estimates. —

3.3.1. Lemma. — *Let $\mathcal{X}$ be a formal R-scheme whose generic fibre $\mathcal{X}_K$ is of dimension $\leq d$. Then*

1. *For every n in $\mathbf{N}$,*
$$\dim \pi_n(\mathrm{Gr}(\mathcal{X})) \leq (n+1)d.$$
2. *For $m \geq n$, the fibres of the projection $\pi_m(\mathrm{Gr}(\mathcal{X})) \to \pi_n(\mathrm{Gr}(\mathcal{X}))$ are of dimension $\leq (m-n)d$.*

3.3.2. Lemma. — *Let $\mathcal{X}$ be a formal R-scheme whose generic fibre $\mathcal{X}_K$ is of dimension d. Let $\mathcal{S}$ be a closed formal R-subscheme of $\mathcal{X}$ such that $\mathcal{S}_K$ is of dimension $< d$. Then, for all integers n, i and ℓ such that $n \geq i \geq \gamma_{\mathcal{S}}(\ell)$, where $\gamma_{\mathcal{S}}$ is the Greenberg function of $\mathcal{S}$ defined in 2.5.2, $\pi_{n,\mathcal{X}}(\pi_{i,\mathcal{X}}^{-1}\mathrm{Gr}_i(\mathcal{S}))$ is of dimension $\leq (n+1)d - \ell - 1$.*

3.4. Grothendieck rings. —

Let k be a field. We denote by $K_0(\mathrm{Var}_k)$ the abelian group generated by symbols $[S]$, for S a variety over k, with the relations $[S] = [S']$ if S and S' are isomorphic and $[S] = [S'] + [S \setminus S']$ if S' is closed in S. There is a natural ring structure on $K_0(\mathrm{Var}_k)$, the product being induced by the cartesian product of varieties, and to any constructible set S in some variety one naturally associates a class $[S]$ in $K_0(\mathrm{Var}_k)$. We denote by $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$ the localisation $K_0(\mathrm{Var}_k)_{\mathrm{loc}} := K_0(\mathrm{Var}_k)[\mathbf{L}^{-1}]$ with $\mathbf{L} := [\mathbf{A}_k^1]$. We denote by $F^m K_0(\mathrm{Var}_k)_{\mathrm{loc}}$ the subgroup generated by $[S]\mathbf{L}^{-i}$ with $\dim S - i \leq -m$, and by $\widehat{K_0(\mathrm{Var}_k)}$ the completion of $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$

with respect to the filtration $F^{\cdot}$[4]. We will also denote by $F^{\cdot}$ the filtration induced on $\widehat{K_0(\mathrm{Var}_k)}$. We denote by $\overline{K_0(\mathrm{Var}_k)_{\mathrm{loc}}}$ the image of $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$ in $\widehat{K_0(\mathrm{Var}_k)}$. We put on the ring $\widehat{K_0(\mathrm{Var}_k)}$ a structure of non-archimedean ring by setting $||a|| := 2^{-n}$, where n is the largest n such that $a \in F^n\widehat{K_0(\mathrm{Var}_k)}$, for $a \neq 0$ and $||0|| = 0$.

3.5. Cylinders. — Let $\mathcal{X}$ be a formal R-scheme. A subset A of $\mathrm{Gr}(\mathcal{X})$ is cylindrical of level $n \geq 0$ if $A = \pi_n^{-1}(C)$ with C a constructible subset of $\mathrm{Gr}_n(\mathcal{X})$. We denote by $\mathbf{C}_{\mathcal{X}}$ the set of cylindrical subsets of $\mathrm{Gr}(\mathcal{X})$ of some level. Let us remark that $\mathbf{C}_{\mathcal{X}}$ is a boolean algebra, *i.e.* contains $\mathrm{Gr}(\mathcal{X})$, $\emptyset$, and is stable by finite intersection, finite union, and by taking complements. It follows from Proposition 3.1.1, that if A is cylindrical of some level, then $\pi_n(A)$ is constructible for every $n \geq 0$.
A basic finiteness property of cylinders is the following:

3.5.1. Lemma. — *Let A_i, $i \in I$, be an enumerable family of cylindrical subsets of* $\mathrm{Gr}(\mathcal{X})$*. If $A := \cup_{i \in I} A_i$ is also a cylinder, then there exists a finite subset J of I such that $A := \cup_{i \in J} A_i$.*

Proof. — Since $\mathrm{Gr}(\mathcal{X})$ is quasi-compact, this follows from Théorème 7.2.5 of [**13**]. □

3.6. Motivic measure for cylinders. — Let $\mathcal{X}$ a flat formal R-scheme of relative dimension d. Let A be a cylinder of $\mathrm{Gr}(\mathcal{X})$. We shall say A is stable of level n if it is cylindrical of level n and if, for every $m \geq n$, the morphism

$$\pi_{m+1}(\mathrm{Gr}(\mathcal{X})) \longrightarrow \pi_m(\mathrm{Gr}(\mathcal{X}))$$

is a piecewise trivial fibration over $\pi_n(A)$ with fibre $\mathbf{A}_k^d$. We denote by $\mathbf{C}_{0,\mathcal{X}}$ the set of stable cylindrical subsets of $\mathrm{Gr}(\mathcal{X})$ of some level.
It follows from Proposition 3.1.2 that every cylinder in $\mathrm{Gr}(\mathcal{X})$ is stable when $\mathcal{X}$ is smooth. When $\mathcal{X}$ is no longer assumed to be smooth, $\mathbf{C}_{0,\mathcal{X}}$ is in general not a boolean algebra, but is an ideal of $\mathbf{C}_{\mathcal{X}}$: $\mathbf{C}_{0,\mathcal{X}}$ contains $\emptyset$, is stable by finite union, and the intersection of an element in $\mathbf{C}_{\mathcal{X}}$ with an element of $\mathbf{C}_{0,\mathcal{X}}$ belongs to $\mathbf{C}_{0,\mathcal{X}}$.
In general $\mathrm{Gr}(\mathcal{X})$ is not stable, but it follows from Proposition 3.2.1 that $\mathrm{Gr}^{(e)}(\mathcal{X})$ is a stable cylinder of $\mathrm{Gr}(\mathcal{X})$, for every $e \geq 0$.
From first principles, one proves (cf. [**10**], [**24**]):

3.6.1. Proposition-Definition. — There is a unique additive morphism

$$\tilde{\mu} : \mathbf{C}_{0,\mathcal{X}} \longrightarrow K_0(\mathrm{Var}_k)_{\mathrm{loc}}$$

such that $\tilde{\mu}(A) = [\pi_n(A)]\,\mathbf{L}^{-(n+1)d}$, when A is a stable cylinder of level n. Furthermore the measure $\tilde{\mu}$ is σ-additive.

One deduces from Proposition 3.3.1 and Proposition 3.3.2, cf. [**10**], [**24**], the following:

---

[4] It is still unknown whether the filtration $F^{\cdot}$ is separated or not.

3.6.2. Proposition. —

1. *For any cylinder* A *in* $\mathbf{C}_{\mathcal{X}}$*, the limit*
$$\mu(A) := \lim_{e \to \infty} \tilde{\mu}(A \cap \mathrm{Gr}^{(e)}(\mathcal{X}))$$
exists in $\widehat{K_0(\mathrm{Var}_k)}$*.*
2. *If* A *belongs to* $\mathbf{C}_{0,\mathcal{X}}$*, then* $\mu(A)$ *coincide with the image of* $\tilde{\mu}(A)$ *in* $\widehat{K_0(\mathrm{Var}_k)}$*.*
3. *The measure* $A \mapsto \mu(A)$ *is* σ*-additive.*
4. *For* A *and* B *in* $\mathbf{C}_{\mathcal{X}}$*,* $||\mu(A \cup B)|| \leq \max(||\mu(A)||, ||\mu(B)||)$*. If* $A \subset B$*,* $||\mu(A)|| \leq ||\mu(B)||$*.*

3.7. Measurable subsets of $\mathrm{Gr}(\mathcal{X})$**.** — For A and B subsets of the same set, we use the notation $A \triangle B$ for $A \cup B \setminus A \cap B$.

3.7.1. Definition. — We say that a subset A of $\mathrm{Gr}(\mathcal{X})$ is *measurable* if, for every positive real number ε, there exists an ε-cylindrical approximation, *i.e.* a sequence of cylindrical subsets $A_i(\varepsilon)$, $i \in \mathbf{N}$, such that
$$\Big(A \triangle A_0(\varepsilon)\Big) \subset \bigcup_{i \geq 1} A_i(\varepsilon),$$
and $||\mu(A_i(\varepsilon))|| \leq \varepsilon$ for all $i \geq 1$. We say that A is *strongly measurable* if moreover we can take $A_0(\varepsilon) \subset A$.

3.7.2. Theorem. — *If* A *is a measurable subset of* $\mathrm{Gr}(\mathcal{X})$*, then*
$$\mu(A) := \lim_{\varepsilon \to 0} \mu(A_0(\varepsilon))$$
exists in $\widehat{K_0(\mathrm{Var}_k)}$ *and is independent of the choice of the sequences* $A_i(\varepsilon)$*,* $i \in \mathbf{N}$*.*

For A a measurable subset of $\mathrm{Gr}(\mathcal{X})$, we shall call $\mu(A)$ the motivic measure of A. We denote by $\mathbf{D}_{\mathcal{X}}$ the set of measurable subsets of $\mathrm{Gr}(\mathcal{X})$.
One should remark that obviously $\mathbf{C}_{\mathcal{X}}$ is contained in $\mathbf{D}_{\mathcal{X}}$.

3.7.3. Proposition. —

1. $\mathbf{D}_{\mathcal{X}}$ *is a boolean algebra.*
2. *If* A_i*,* $i \in \mathbf{N}$*, is a sequence of measurable subsets of* $\mathrm{Gr}(\mathcal{X})$ *with* $\lim_{i \to \infty} ||\mu(A_i)|| = 0$*, then* $\cup_{i \in \mathbf{N}} A_i$ *is measurable.*
3. *Let* A_i*,* $i \in \mathbf{N}$*, be a family of measurable subsets of* $\mathrm{Gr}(\mathcal{X})$*. Assume the sets* A_i *are mutually disjoint and that* $A := \cup_{i \in \mathbf{N}} A_i$ *is measurable. Then* $\sum_{i \in \mathbf{N}} \mu(A_i)$ *converges in* $\widehat{K_0(\mathrm{Var}_k)}$ *to* $\mu(A)$*.*
4. *If* A *and* B *are measurable subsets of* $\mathrm{Gr}(\mathcal{X})$ *and if* $A \subset B$*, then* $||\mu(A)|| \leq ||\mu(B)||$*.*

3.7.4. Remark. — In the situation of Remark 2.3.3, one can check that the notion of cylinders, stable cylinders, measurable subsets of $\mathrm{Gr}(X \widehat{\otimes} R)$ coincides with the analogous notions introduced in [**11**] for subsets of $\mathcal{L}(X)$.

3.8. Order of the Jacobian ideal. — Let $h : \mathcal{Y} \to \mathcal{X}$ be a morphism of flat formal R-schemes of relative dimension d.
Let y be a point of $\mathrm{Gr}(\mathcal{Y}) \setminus \mathrm{Gr}(\mathcal{Y}_{\mathrm{sing}})$ defined over some field extension F of k. We denote by $\varphi : \mathrm{Spf}\, R_F \to \mathcal{Y}$ the corresponding morphism of formal R-schemes. We define $\mathrm{ord}_{\varpi}(\mathrm{Jac}_h)(y)$, the order of the Jacobian ideal of h at y, as follows.
From the natural morphism $h^*\Omega_{\mathcal{X}|R} \to \Omega_{\mathcal{Y}|R}$, one deduces, by taking the d-th exterior power, a morphism $h^*\Omega^d_{\mathcal{X}|R} \to \Omega^d_{\mathcal{Y}|R}$, hence a morphism

$$(\varphi^* h^* \Omega^d_{\mathcal{X}|R})/(\mathrm{torsion}) \longrightarrow (\varphi^* \Omega^d_{\mathcal{Y}|R})/(\mathrm{torsion}).$$

Since $L := (\varphi^*\Omega^d_{\mathcal{Y}|R})/(\mathrm{torsion})$ is a free $\mathcal{O}_{R_F}$-module of rank 1, it follows from the structure theorem for finite type modules over principal domains, that the image of $M := (\varphi^* h^* \Omega^d_{\mathcal{X}|R})/(\mathrm{torsion})$ in L is either 0, in which case we set $\mathrm{ord}_{\varpi}(\mathrm{Jac}_h)(y) = \infty$, or $\varpi^n L$, for some $n \in \mathbf{N}$, in which case we set $\mathrm{ord}_{\varpi}(\mathrm{Jac}_h)(y) = n$.

3.9. The change of variable formula. — If $h : \mathcal{Y} \to \mathcal{X}$ is a morphism of formal R-schemes, we still write h for the corresponding morphism $\mathrm{Gr}(\mathcal{Y}) \to \mathrm{Gr}(\mathcal{X})$.

The following lemmas are basic geometric ingredients in the proof of the change of variable formula 3.9.3 and 3.9.4.

3.9.1. Lemma. — *Let $h : \mathcal{Y} \to \mathcal{X}$ be a morphism between flat formal R-schemes of relative dimension d. We assume $\mathcal{Y}$ is smooth. For e and e' in $\mathbf{N}$ we set*

$$\Delta_{e,e'} := \Big\{\varphi \in \mathrm{Gr}(\mathcal{Y}) \;\Big|\; \mathrm{ord}_{\varpi}(\mathrm{Jac}_h)(y) = e \quad \text{and} \quad h(\varphi) \in \mathrm{Gr}^{(e')}(\mathcal{X})\Big\}.$$

Then, there exists c in $\mathbf{N}$ such that, for every $n \geq 2e$, $n \geq e + ce'$, for every φ in $\Delta_{e,e'}$ and every x in $\mathrm{Gr}(\mathcal{X})$ such that $\pi_n(h(\varphi)) = \pi_n(x)$, there exists y in $\mathrm{Gr}(\mathcal{Y})$ such that $h(y) = x$ and $\pi_{n-e}(\varphi) = \pi_{n-e}(y)$.

3.9.2. Lemma. — *Let $h : \mathcal{Y} \to \mathcal{X}$ be a morphism between flat formal R-schemes of relative dimension d. We assume $\mathcal{Y}$ is smooth. Let B be a cylinder in $\mathrm{Gr}(\mathcal{Y})$ and set $A = h(B)$. Assume $\mathrm{ord}_{\varpi}(\mathrm{Jac}_h)$ is constant with value $e < \infty$ on B and that $A \subset \mathrm{Gr}^{(e')}(\mathcal{X})$ for some $e' \geq 0$. Then A is a cylinder. Furthermore, if the restriction of h to B is injective, then for $n \gg 0$, the following holds:*

1. *If φ and φ' belong to B and $\pi_n(h(\varphi)) = \pi_n(h(\varphi'))$, then $\pi_{n-e}(\varphi) = \pi_{n-e}(\varphi')$.*
2. *The morphism $\pi_n(B) \to \pi_n(A)$ induced by h is a piecewise trivial fibration with fibre $\mathbf{A}^e_k$.*

For a measurable subset A of $\mathrm{Gr}(\mathcal{X})$ and a function $\alpha : A \to \mathbf{Z} \cup \{\infty\}$, we say that $\mathbf{L}^{-\alpha}$ is integrable or that α is exponentially integrable if the fibres of α are measurable and if the motivic integral

$$\int_A \mathbf{L}^{-\alpha} d\mu := \sum_{n \in \mathbf{Z}} \mu(A \cap \alpha^{-1}(n))\mathbf{L}^{-n}$$

converges in $\widehat{K_0(\mathrm{Var}_k)}$.

When all the fibres $A \cap \alpha^{-1}(n)$ are stable cylinders and α takes only a finite number of values on A, it is not necessary to go to the completion of $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$ and one may define directly

$$\int_A \mathbf{L}^{-\alpha} d\tilde{\mu} := \sum_{n \in \mathbf{Z}} \tilde{\mu}(A \cap \alpha^{-1}(n)) \mathbf{L}^{-n}$$

in $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$.

3.9.3. Theorem. — *Let $h : \mathcal{Y} \to \mathcal{X}$ be a morphism between flat formal R-schemes of relative dimension d. We assume $\mathcal{Y}$ is smooth. Let B be a strongly measurable subset of $\mathrm{Gr}(\mathcal{Y})$. Assume h induces a bijection between B and $A := h(B)$. Then, for every exponentially integrable function $\alpha : A \to \mathbf{Z} \cup \infty$, the function $\alpha \circ h + \mathrm{ord}_{\varpi}(\mathrm{Jac}_h)$ is exponentially integrable on B and*

$$\int_A \mathbf{L}^{-\alpha} d\mu = \int_B \mathbf{L}^{-\alpha \circ h - \mathrm{ord}_{\varpi}(\mathrm{Jac}_h)} d\mu.$$

We shall also need the following variant of Theorem 3.9.3.

3.9.4. Theorem. — *Let $h : \mathcal{Y} \to \mathcal{X}$ be a morphism between flat formal R-schemes of relative dimension d. We assume $\mathcal{Y}$ and $\mathcal{X}_K$ are smooth and that the morphism $h_K : \mathcal{Y}_K \to \mathcal{X}_K$ induced by h is étale (cf. [**8**]). Let B be a cylinder in $\mathrm{Gr}(\mathcal{Y})$. Assume h induces a bijection between B and $A := h(B)$ and that A is a stable cylinder of $\mathrm{Gr}(\mathcal{X})$. Then the fibres $B \cap \mathrm{ord}_{\varpi}(\mathrm{Jac}_h)^{-1}(n)$ are stable cylinders, $\mathrm{ord}_{\varpi}(\mathrm{Jac}_h)^{-1}(n)$ takes only a finite number of values on B, and*

$$\int_A d\tilde{\mu} = \int_B \mathbf{L}^{-\mathrm{ord}_{\varpi}(\mathrm{Jac}_h)} d\tilde{\mu}$$

in $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$.

4. Integration on smooth rigid varieties

4.1. Order of differential forms. — Let $\mathcal{X}$ be a flat formal R-scheme equidimensional of relative dimension d. Consider a differential form ω in $\Omega^d_{\mathcal{X}|R}(\mathcal{X})$. Let x be a point of $\mathrm{Gr}(\mathcal{X}) \setminus \mathrm{Gr}(\mathcal{X}_{\mathrm{sing}})$ defined over some field extension F of k. We denote by $\varphi : \mathrm{Spf}\, R_F \to \mathcal{X}$ the corresponding morphism of formal R-schemes. Since $L := (\varphi^* \Omega^d_{\mathcal{X}|R})/(\mathrm{torsion})$ is a free $\mathcal{O}_{R_F}$-module of rank 1, it follows from the structure theorem for finite type modules over principal domains, that its submodule M generated by $\varphi^* \omega$ is either 0, in which case we set $\mathrm{ord}_{\varpi}(\omega)(x) = \infty$, or $\varpi^n L$, for some $n \in \mathbf{N}$, in which case we set $\mathrm{ord}_{\varpi}(\omega)(x) = n$.

Since there is a canonical isomorphism $\Omega^d_{\mathcal{X}_K}(\mathcal{X}_K) \simeq \Omega^d_{\mathcal{X}|R}(\mathcal{X}) \otimes_R K$ (cf. Proposition 1.5 of [**8**]), if ω is in $\Omega^d_{\mathcal{X}_K}(\mathcal{X}_K)$, we write $\omega = \varpi^{-n} \tilde{\omega}$, with $\tilde{\omega}$ in $\Omega^d_{\mathcal{X}|R}(\mathcal{X})$ and $n \in \mathbf{N}$, we set $\mathrm{ord}_{\varpi, \mathcal{X}}(\omega) := \mathrm{ord}_{\varpi}(\tilde{\omega}) - n$. Clearly this definition does not depend on the choice of $\tilde{\omega}$.

4.1.1. Lemma. — *Let $h : \mathcal{Y} \to \mathcal{X}$ be a morphism between flat formal R-schemes equidimensional of relative dimension d. Let ω be in $\Omega^d_{\mathcal{X}|R}(\mathcal{X})$ (resp. in $\Omega^d_{\mathcal{X}_K}(\mathcal{X}_K)$). Let y be a point in $\mathrm{Gr}(\mathcal{Y}) \setminus \mathrm{Gr}(\mathcal{Y}_{\mathrm{sing}})$ and assume $h(y)$ belongs to $\mathrm{Gr}(\mathcal{X}) \setminus \mathrm{Gr}(\mathcal{X}_{\mathrm{sing}})$. Then*

$$\mathrm{ord}_\varpi(h^*\omega)(y) = \mathrm{ord}_\varpi(\omega)(h(y)) + \mathrm{ord}_\varpi(\mathrm{Jac}_h)(y),$$

resp.

$$\mathrm{ord}_{\varpi,\mathcal{Y}}(h_K^*\omega)(y) = \mathrm{ord}_{\varpi,\mathcal{X}}(\omega)(h(y)) + \mathrm{ord}_\varpi(\mathrm{Jac}_h)(y).$$

Proof. — Follows directly from the definitions. □

4.1.2. Theorem-Definition. — Let X be a smooth rigid variety over K, purely of dimension d. Let ω be a differential form in $\Omega^d_X(X)$.

(1) Let $\mathcal{X}$ be a formal R-model of X. Then the function $\mathrm{ord}_{\varpi,\mathcal{X}}(\omega)$ is exponentially integrable on $\mathrm{Gr}(\mathcal{X})$ and the integral $\int_{\mathrm{Gr}(\mathcal{X})} \mathbf{L}^{-\mathrm{ord}_{\varpi,\mathcal{X}}(\omega)} d\mu$ in $\widehat{K_0(\mathrm{Var}_k)}$ does not depend on the model $\mathcal{X}$. We denote it by $\int_X \omega d\mu$.

(2) Assume furthermore ω is a gauge form, *i.e.* that it generates Ω^d_X at every point of X, and assume some open dense formal subscheme $\mathcal{U}$ of $\mathcal{X}$ is a weak Néron model of X. Then the function $\mathrm{ord}_{\varpi,\mathcal{X}}(\omega)$ takes only a finite number of values and its fibres are stable cylinders. Furthermore the integral $\int_{\mathrm{Gr}(\mathcal{X})} \mathbf{L}^{-\mathrm{ord}_{\varpi,\mathcal{X}}(\omega)} d\tilde{\mu}$ in $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$ does not depend on the model $\mathcal{X}$. We denote it by $\int_X \omega d\tilde{\mu}$.

Proof. — Let us prove (2). Write $\omega = \varpi^{-n}\tilde{\omega}$, with $\tilde{\omega}$ in $\Omega^d_{\mathcal{X}|R}(\mathcal{X})$ and $n \in \mathbf{N}$. Since $\mathcal{U}$ is smooth, $\Omega^d_{\mathcal{U}|R}$ is locally free of rank 1 and $\tilde{\omega}\mathcal{O}_U \otimes (\Omega^d_{\mathcal{U}|R})^{-1}$ is isomorphic to a principal ideal sheaf $(f)\mathcal{O}_U$, with f in $\mathcal{O}_U$. Furthermore, the function $\mathrm{ord}_{\varpi,\mathcal{X}}(\tilde{\omega})$ coincides with the function $\mathrm{ord}_\varpi(f)$ which to a point φ of $\mathrm{Gr}(\mathcal{U}) = \mathrm{Gr}(\mathcal{X})$ associates $\mathrm{ord}_\varpi(f(\varphi))$. The fibres of $\mathrm{ord}_\varpi(f)$ are stable cylinders. Since ω is a gauge form, f induces an invertible function on X, hence, by the maximum principle (cf. [**4**]), the function $\mathrm{ord}_\varpi(f)$ takes only a finite number of values. To prove that $\int_{\mathrm{Gr}(\mathcal{X})} \mathbf{L}^{-\mathrm{ord}_{\varpi,\mathcal{X}}(\omega)} d\tilde{\mu}$ in $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$ does not depend on the model $\mathcal{X}$, it is enough to consider the case of another model $\mathcal{X}'$ obtained from $\mathcal{X}$ by an admissible formal blow-up $h : \mathcal{X}' \to \mathcal{X}$. We may also assume $\mathcal{X}'$ contains as an open dense formal subscheme a weak Néron model $\mathcal{U}'$ of X. The equality

$$\int_{\mathrm{Gr}(\mathcal{X}')} \mathbf{L}^{-\mathrm{ord}_{\varpi,\mathcal{X}'}(\omega)} d\tilde{\mu} = \int_{\mathrm{Gr}(\mathcal{X})} \mathbf{L}^{-\mathrm{ord}_{\varpi,\mathcal{X}}(\omega)} d\tilde{\mu}$$

then follows from Lemma 4.1.1 and Theorem 3.9.4. Statement (1) follows similarly from Lemma 4.1.1 and Theorem 3.9.3. □

4.1.3. Remark. — A situation where gauge forms naturally occur is that of reductive groups. Let G be a connected reductive group over k. B. Gross constructs in [**16**], using Bruhat-Tits theory, a differential form of top degree ω_G on G which is defined up to multiplication by a unit in R. One may easily check that the differential form ω_G induces a gauge form on the rigid K-group $G^{\mathrm{rig}} := (G\widehat{\otimes}R)_K$.

4.1.4. Lemma. — *Let X be a smooth rigid variety over K, purely of dimension d, and let ω be a gauge form on X. Let $\mathcal{O} = (O_i)_{i \in J}$ be a finite admissible covering and set $O_I := \cap_{i \in I} O_i$ for $I \subset J$. Then*

$$\int_X \omega d\tilde{\mu} = \sum_{\emptyset \neq I \subset J} (-1)^{|I|-1} \int_{O_I} \omega_{|O_I} d\tilde{\mu}.$$

If ω is only assumed to be a differential form in $\Omega^d_X(X)$, then

$$\int_X \omega d\mu = \sum_{\emptyset \neq I \subset J} (-1)^{|I|-1} \int_{O_I} \omega_{|O_I} d\mu.$$

Proof. — Let us prove the first statement, the proof of the second one being similar. It is enough to consider the case $|J| = 2$. Choose an R-model $\mathcal{X}$ containing a weak Néron model $\mathcal{U}$ of X as an open dense formal subscheme and such that the covering $X = O_1 \cup O_2$ is induced from a covering $\mathcal{X} = \mathcal{O}_1 \cup \mathcal{O}_2$ by open formal subschemes. It is sufficient to prove that

$$\int_{\mathrm{Gr}(\mathcal{X})} \mathbf{L}^{-\mathrm{ord}_{\varpi,\mathcal{X}}(\omega)} d\tilde{\mu} = \int_{\mathrm{Gr}(\mathcal{O}_1)} \mathbf{L}^{-\mathrm{ord}_{\varpi,\mathcal{O}_1}(\omega_{|O_1})} d\tilde{\mu} + \int_{\mathrm{Gr}(\mathcal{O}_2)} \mathbf{L}^{-\mathrm{ord}_{\varpi,\mathcal{O}_2}(\omega_{|O_2})} d\tilde{\mu}$$
$$- \int_{\mathrm{Gr}(\mathcal{O}_1 \cap \mathcal{O}_2)} \mathbf{L}^{-\mathrm{ord}_{\varpi,\mathcal{O}_1 \cap \mathcal{O}_2}(\omega_{|O_1 \cap O_2})} d\tilde{\mu},$$

which follows from the fact that for every open formal subscheme $\mathcal{O}$ of $\mathcal{X}$ the function $\mathrm{ord}_{\varpi,\mathcal{X}}(\omega)$ restricts to $\mathrm{ord}_{\varpi,\mathcal{O}}(\omega_{|\mathcal{O}_K})$ on $\mathrm{Gr}(\mathcal{O})$ and the equalities $\mathrm{Gr}(\mathcal{X}) = \mathrm{Gr}(\mathcal{O}_1) \cup \mathrm{Gr}(\mathcal{O}_2)$ and $\mathrm{Gr}(\mathcal{O}_1) \cap \mathrm{Gr}(\mathcal{O}_2) = \mathrm{Gr}(\mathcal{O}_1 \cap \mathcal{O}_2)$ which follow from Proposition 2.3.2. □

4.1.5. Proposition. — *Let X and X' be smooth rigid K-varieties purely of dimension d and d' and let ω and ω' be gauge forms on X and X'. Then*

$$\int_{X \times X'} \omega \times \omega' d\tilde{\mu} = \int_X \omega d\tilde{\mu} \times \int_{X'} \omega' d\tilde{\mu}.$$

If ω and ω' are only assumed to be differential forms in $\Omega^d_X(X)$, then

$$\int_{X \times X'} \omega \times \omega' d\mu = \int_X \omega d\mu \times \int_{X'} \omega' d\mu.$$

Proof. — Let us prove the first assertion, the proof of the second one being similar. Choose R-models $\mathcal{X}$ and $\mathcal{X}'$ of X and X' respectively containing a weak Néron model $\mathcal{U}$ of X and $\mathcal{U}'$ of X' as an open dense formal subscheme. Also write $\omega = \varpi^{-n}\tilde{\omega}$ and $\omega' = \varpi^{-n'}\tilde{\omega}'$, with $\tilde{\omega}$ and $\tilde{\omega}'$ in $\Omega^d_{\mathcal{X}|R}(\mathcal{X})$ and $\Omega^{d'}_{\mathcal{X}'|R}(\mathcal{X}')$, respectively. It enough to check that $\tilde{\mu}(\mathrm{ord}_{\varpi,\mathcal{X}\times\mathcal{X}'}(\tilde{\omega} \times \tilde{\omega}') = m)$ is equal to $\sum_{m'+m''=m} \tilde{\mu}(\mathrm{ord}_{\varpi,\mathcal{X}}(\tilde{\omega}) = m') \times \tilde{\mu}(\mathrm{ord}_{\varpi,\mathcal{X}'}(\tilde{\omega}') = m'')$, which follows the fact that on $\mathrm{Gr}(\mathcal{X} \times \mathcal{X}') \simeq \mathrm{Gr}(\mathcal{X}) \times \mathrm{Gr}(\mathcal{X}') = \mathrm{Gr}(\mathcal{U}) \times \mathrm{Gr}(\mathcal{U}')$, the functions $\mathrm{ord}_{\varpi,\mathcal{X}\times\mathcal{X}'}(\tilde{\omega} \times \tilde{\omega}')$ and $\mathrm{ord}_{\varpi,\mathcal{X}}(\tilde{\omega}) + \mathrm{ord}_{\varpi,\mathcal{X}'}(\tilde{\omega}')$ are equal. □

4.2. Invariants for gauged smooth rigid varieties. — Let d be an integer ≥ 0. We define $K_0(\mathrm{GSRig}_K^d)$, the Grothendieck group of gauged smooth rigid K-varieties of dimension d, as follows: as an abelian group it is the quotient of the free abelian group over symbols $[X, \omega]$, with X a smooth rigid K-variety of dimension d and ω a gauge form on X by the relations

$$[X', \omega'] = [X, \omega]$$

if there is an isomorphism $h : X' \to X$ with $h^*\omega = \omega'$, and

$$[X, \omega] = \sum_{\emptyset \neq I \subset J} (-1)^{|I|-1} [O_I, \omega_{|O_I}],$$

when $(O_i)_{i \in J}$ is a finite admissible covering of X, with the notation $O_I := \cap_{i \in I} O_i$ for $I \subset J$. One puts a graded ring structure on $K_0(\mathrm{GSRig}_K) := \oplus_d K_0(\mathrm{GSRig}_K^d)$ by requiring

$$[X, \omega] \times [X', \omega'] := [X \times X', \omega \times \omega'].$$

Forgetting gauge forms, one defines similarly $K_0(\mathrm{SRig}_K^d)$, the Grothendieck group of smooth rigid K-varieties of dimension d, and the graded ring $K_0(\mathrm{SRig}_K) := \oplus_d K_0(\mathrm{SRig}_K^d)$. There are natural forgetful morphisms

$$F : K_0(\mathrm{GSRig}_K^d) \longrightarrow K_0(\mathrm{SRig}_K^d)$$

and

$$F : K_0(\mathrm{GSRig}_K) \longrightarrow K_0(\mathrm{SRig}_K).$$

4.2.1. ***Proposition.*** — *The assignment which to a gauged smooth rigid K-variety (X, ω) associates $\int_X \omega d\tilde{\mu}$ factorizes uniquely as a ring morphism*

$$\tilde{\mu} : K_0(\mathrm{GSRig}_K) \to K_0(\mathrm{Var}_k)_{\mathrm{loc}}.$$

Proof. — This follows from Lemma 4.1.4 and Proposition 4.1.5. □

4.3. A formula for $\int_X \omega d\tilde{\mu}$. — Let X be a smooth rigid variety over K of pure dimension d. Let $\mathcal{U}$ be a weak Néron model of X contained in some model $\mathcal{X}$ of X and let ω be a form in $\Omega^d_{\mathcal{X}|R}(\mathcal{X})$ inducing a gauge form on X. We denote by U_0^i, $i \in J$, the irreducible components of the special fibre of $\mathcal{U}$. By assumption, each U_0^i is smooth and $U_0^i \cap U_0^j = \emptyset$ for $i \neq j$. We denote by $\mathrm{ord}_{U_0^i}(\omega)$ the unique integer n such that $\varpi^{-n}\omega$ generates $\Omega^d_{\mathcal{X}|R}$ at the generic point of U_0^i. More generally, if ω is a gauge form in $\Omega^d_{\mathcal{X}_K}(\mathcal{X}_K)$, we write $\omega = \varpi^{-n}\tilde{\omega}$, with $\tilde{\omega}$ in $\Omega^d_{\mathcal{X}|R}(\mathcal{X})$ and $n \in \mathbf{N}$, and we set $\mathrm{ord}_{U_0^i}(\omega) := \mathrm{ord}_{U_0^i}(\tilde{\omega}) - n$.

4.3.1. ***Proposition.*** — *Let X be a smooth rigid variety over K of pure dimension d. Let $\mathcal{U}$ be a weak Néron model of X contained in some model $\mathcal{X}$ of X and let ω be a gauge form in $\Omega^d_{\mathcal{X}_K}(\mathcal{X}_K)$. With the above notations, we have*

$$\int_X \omega d\tilde{\mu} = \mathbf{L}^{-d} \sum_{i \in J} [U_0^i] \, \mathbf{L}^{-\mathrm{ord}_{U_0^i}(\omega)}$$

in $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$.

Proof. — Denote by $\mathcal{U}_0^i$ the irreducible component of $\mathcal{X}$ with special fibre U_0^i. Since $\mathrm{Gr}(\mathcal{X})$ is the disjoint union of the sets $\mathrm{Gr}(\mathcal{U}_0^i)$, we may assume $\mathcal{X}$ is a smooth irreducible formal R-scheme of dimension d. Let ω be a section of $\Omega^d_{\mathcal{X}|R}(\mathcal{X})$ which generates $\Omega^d_{\mathcal{X}|R}$ at the generic point of $\mathcal{X}$ and induces a gauge form on the generic fibre. Let us remark that the function $\mathrm{ord}_{\varpi,\mathcal{X}}(\omega)$ is identically equal to 1 on $\mathrm{Gr}(\mathcal{X})$. Indeed, after shrinking $\mathcal{X}$, we may write $\omega = f\omega_0$ with ω_0 a generator $\Omega^d_{\mathcal{X}|R}$ a every point and f in $\mathcal{O}_{\mathcal{X}}(\mathcal{X})$. By hypothesis f is a unit at the generic point of $\mathcal{X}$. Assume at some point x of $\mathrm{Gr}(\mathcal{X})$, $\mathrm{ord}_{\varpi} f(x) \geq 1$; it would follow that the locus of $f = 0$ is non empty in $\mathcal{X}$, which contradicts the assumption that ω induces a gauge form on the generic fibre. Hence, we get $\int_{\mathrm{Gr}(\mathcal{X})} \mathbf{L}^{-\mathrm{ord}_{\varpi,\mathcal{X}}(\omega)} \tilde{d}\mu = \mathbf{L}^{-d}[X_0]$, and the result follows. □

4.4. Application to Calabi-Yau varieties over K. — Let X be a Calabi-Yau variety over K. By this we mean a smooth projective algebraic variety over K of pure dimension d with Ω^d_X trivial. We denote by X^{an} the rigid K-variety associated to X. Since X is proper, X^{an} is canonically isomorphic to the generic fibre of the formal completion $X\widehat{\otimes}R$. By GAGA (cf. [**20**] Theorem 2.8), $\Omega^d_{X^{\mathrm{an}}}(X^{\mathrm{an}}) = \Omega^d_X(X) = K$. Now we can associate to any Calabi-Yau variety over K a canonical element in the ring $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$ which coincides with the class of the special fibre when X has a model with good reduction.

4.4.1. ***Theorem.*** — *Let X be a Calabi-Yau variety over K, let $\mathcal{U}$ be a weak Néron model of X^{an} and let ω be a gauge form on X^{an}. We denote by U_0^i, $i \in J$, the irreducible components of the special fibre of $\mathcal{U}$ and set $\alpha(\omega) := \inf \mathrm{ord}_{U_0^i}(\omega)$. Then the virtual variety*

$$[\overline{X}] := \sum_{i\in J} [U_0^i]\, \mathbf{L}^{\alpha(\omega) - \mathrm{ord}_{U_0^i}(\omega)} \tag{4.4.1}$$

in $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$ only depends on X. When X has a proper smooth model with good reduction over R, $[\overline{X}]$ is equal to the class of the special fibre.

Proof. — Let ω be a gauge form on X^{an}. By Proposition 4.3.1, the right hand side of (4.4.1) is equal to $\mathbf{L}^{d-\alpha(\omega)} \int_{X^{an}} \omega d\tilde{\mu}$ which does not depend on ω. □

In particular, we have the following analogue of Batyrev's result on birational projective Calabi-Yau manifolds [**1**], [**10**].

4.4.2. ***Corollary.*** — *Let X be a Calabi-Yau variety over K and let $\mathcal{X}$ and $\mathcal{X}'$ be two proper and smooth R-models of X with special fibres $\mathcal{X}_0$ and $\mathcal{X}'_0$. Then*

$$[\mathcal{X}_0] = [\mathcal{X}'_0]$$

in $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$. □

4.4.3. Remark. — Calabi-Yau varieties over $k((t))$, with k of characteristic zero have been considered in [**18**].

4.5. A motivic Serre invariant for smooth rigid varieties. — We can now define the motivic Serre invariant for smooth rigid varieties.

4.5.1. Theorem. — *There is a canonical ring morphism*

$$\lambda : K_0(\mathrm{SRig}_K) \longrightarrow K_0(\mathrm{Var}_k)_{\mathrm{loc}}/(\mathbf{L}-1)K_0(\mathrm{Var}_k)_{\mathrm{loc}}$$

such that the diagram

$$\begin{tikzcd}
K_0(\mathrm{GSRig}_K) \arrow[r, "\tilde{\mu}"] \arrow[d, "F"] & K_0(\mathrm{Var}_k)_{\mathrm{loc}} \arrow[d] \\
K_0(\mathrm{SRig}_K) \arrow[r, "\lambda"] & K_0(\mathrm{Var}_k)_{\mathrm{loc}}/(\mathbf{L}-1)K_0(\mathrm{Var}_k)_{\mathrm{loc}}
\end{tikzcd}$$

is commutative.

Proof. — Since any smooth rigid K-variety of dimension d admits a finite admissible covering by affinoids $(O_i)_{i\in J}$, with $\Omega^d_{O_i}$ trivial, the morphism F is surjective. Hence it is enough to show the following statement: let $\mathcal{X}$ be a smooth formal R-scheme of relative dimension d with $\Omega^d_{\mathcal{X}|R}$ trivial, and let ω_1 and ω_2 be two global sections of $\Omega^d_{\mathcal{X}|R}$ inducing gauge forms on the generic fibre $\mathcal{X}_K$, then $\int_{\mathrm{Gr}(\mathcal{X})}(\mathbf{L}^{-\mathrm{ord}_{\varpi,\mathcal{X}}(\omega_1)} - L^{-\mathrm{ord}_{\varpi,\mathcal{X}}(\omega_2)})d\tilde{\mu}$ belongs to $(\mathbf{L}-1)K_0(\mathrm{Var}_k)_{\mathrm{loc}}$. To prove this, we take ω_0 a global section of $\Omega^d_{\mathcal{X}|R}$ such that $\Omega^d_{\mathcal{X}|R} \simeq \omega_0\mathcal{O}_{\mathcal{X}}$. If ω is any global section of $\Omega^d_{\mathcal{X}|R}$, write $\omega = f\omega_0$ with f in $\mathcal{O}_{\mathcal{X}}(\mathcal{X})$. By the maximum principle, the function $\mathrm{ord}_{\varpi}(f)$ takes only a finite number of values on $\mathrm{Gr}(\mathcal{X})$. It follows we may write $\mathrm{Gr}(\mathcal{X})$ as a disjoint union of the subsets $\mathrm{Gr}(\mathcal{X})_{\mathrm{ord}_{\varpi}(f)=n}$ where $\mathrm{ord}_{\varpi}(f)$ takes the value n. These subsets are stable cylinders and only a finite number of them are non empty. Hence the equality

$$\int_{\mathrm{Gr}(\mathcal{X})}(\mathbf{L}^{-\mathrm{ord}_{\varpi,\mathcal{X}}(\omega)} - L^{-\mathrm{ord}_{\varpi,\mathcal{X}}(\omega_0)})d\tilde{\mu} = \sum_n(\mathbf{L}^{-n}-1)\tilde{\mu}(\mathrm{Gr}(\mathcal{X})_{\mathrm{ord}_{\varpi}(f)=n})$$

holds in $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$ and the statement follows. □

4.5.2. Remark. — The ring $K_0(\mathrm{Var}_k)_{\mathrm{loc}}/(\mathbf{L}-1)K_0(\mathrm{Var}_k)_{\mathrm{loc}}$ is much smaller than $K_0(\mathrm{Var}_k)_{\mathrm{loc}}$, but still quite large. Let ℓ be a prime number distinct from the characteristic of k. Then the étale ℓ-adic Euler characteristic with compact supports $X \mapsto \chi_{c,\ell}(X) := \sum(-1)^i \dim H^i_{c,\text{ét}}(X, \mathbf{Q}_\ell)$ induces a ring morphism $\chi_{c,\ell} : K_0(\mathrm{Var}_k)_{\mathrm{loc}}/(\mathbf{L}-1)K_0(\mathrm{Var}_k)_{\mathrm{loc}} \to \mathbf{Z}$. Similarly, assume there is a natural morphism $H : K_0(\mathrm{Var}_k)_{\mathrm{loc}} \to \mathbf{Z}[u,v]$ which to the class of a variety X over k assigns its Hodge polynomial $H(X)$ for de Rham cohomology with compact support. Such a morphism is known to exist when k is of characteristic zero. Then if one sets $H_{1/2}(X)(u) := H(X)(u, u^{-1})$ one gets a morphism $H_{1/2} : K_0(\mathrm{Var}_k)_{\mathrm{loc}}/(\mathbf{L}-1)K_0(\mathrm{Var}_k)_{\mathrm{loc}} \to \mathbf{Z}[u]$, since $H(\mathbf{A}^1_k) = uv$.

4.5.3. ***Theorem***. — *Let X be a smooth rigid variety over K of pure dimension d. Let $\mathcal{U}$ be a weak Néron model of X and denote by U_0 its special fibre. Then*

$$\lambda([X]) = [U_0]$$

in $K_0(\mathrm{Var}_k)_{\mathrm{loc}}/(\mathbf{L}-1)K_0(\mathrm{Var}_k)_{\mathrm{loc}}$.
In particular, the class of $[U_0]$ *in* $K_0(\mathrm{Var}_k)_{\mathrm{loc}}/(\mathbf{L}-1)K_0(\mathrm{Var}_k)_{\mathrm{loc}}$ *does not depend on the weak Néron model* $\mathcal{U}$.

Proof. — By taking an appropriate admissible cover, we may assume there exists a gauge form on X, in which case the result follows from Proposition 4.3.1, since $[U_0] = \sum_{i\in J}[U_0^i]$. (In fact, one can also prove Theorem 4.5.1 that way, but we prefered to give a proof which is quite parallel to that of Serre in [**25**].) $\square$

4.6. Relation with p-adic integrals on compact locally analytic varieties. — Let K be a local field with residue field $k = \mathbf{F}_q$. Let us consider the Grothendieck group $K_0(\mathrm{SLocAn}_K^d)$ of compact locally analytic smooth varieties over K of pure dimension d, which is defined similarly as $K_0(\mathrm{SRig}_K^d)$ replacing smooth rigid varieties by compact locally analytic smooth varieties and finite admissible covers by finite covers. Also a nowhere vanishing locally analytic d-form on a smooth compact locally analytic variety X of pure dimension d will be called a gauge form on X, and one defines the Grothendieck group $K_0(\mathrm{GSLocAn}_K^d)$ of gauged compact locally analytic smooth varieties over K of pure dimension d similarly as $K_0(\mathrm{GSRig}_K^d)$. There are canonical forgetful morphisms $F : K_0(\mathrm{SRig}_K^d) \to K_0(\mathrm{SLocAn}_K^d)$ and $F : K_0(\mathrm{GSRig}_K^d) \to K_0(\mathrm{GSLocAn}_K^d)$ induced from the functor which to a rigid variety (resp. gauged variety) associates the underlying locally analytic variety (resp. gauged variety). If (X,ω) is a gauged compact locally analytic smooth variety, the p-adic integral $\int_X |\omega|$ belongs to $\mathbf{Z}[q^{-1}]$ (cf. [**25**]), and by additivity of p-adic integrals one gets a morphism $\mathrm{int}_p : K_0(\mathrm{GSLocAn}_K^d) \to \mathbf{Z}[q^{-1}]$.
On the other hand, there is a canonical morphism $N : K_0(\mathrm{Var}_k) \to \mathbf{Z}$ which to the class of a k-variety S assigns the number of points of $S(\mathbf{F}_q)$, and which induces a morphism $N : K_0(\mathrm{Var}_k)_{\mathrm{loc}} \to \mathbf{Z}[q^{-1}]$. We shall also denote by N the induced morphism $K_0(\mathrm{Var}_k)_{\mathrm{loc}}/(\mathbf{L}-1)K_0(\mathrm{Var}_k)_{\mathrm{loc}} \to \mathbf{Z}[q^{-1}]/(q-1)\mathbf{Z}[q^{-1}] \simeq \mathbf{Z}/(q-1)\mathbf{Z}$.

4.6.1. ***Proposition***. — *Let K be a local field with residue field $k = \mathbf{F}_q$. Then the diagram*

$$\begin{tikzcd}
K_0(\mathrm{GSRig}_K^d) \arrow[r, "\tilde{\mu}"] \arrow[d, "F"] & K_0(\mathrm{Var}_k)_{\mathrm{loc}} \arrow[d, "N"] \\
K_0(\mathrm{GSLocAn}_K^d) \arrow[r, "\mathrm{int}_p"] & \mathbf{Z}[q^{-1}]
\end{tikzcd}$$

is commutative.

Proof. — One reduces to showing the following: let $\mathcal{X}$ be a smooth formal R-scheme of dimension d and let f be a function in $\mathcal{O}_{\mathcal{X}}(\mathcal{X})$ which induces a non vanishing function on $\mathcal{X}_K$, then

$$N(\int_{\mathrm{Gr}(\mathcal{X})} \mathbf{L}^{-\mathrm{ord}_\varpi(f)} d\tilde{\mu}) = \int_{\mathcal{X}(R)} q^{-\mathrm{ord}_\varpi(f)} d\tilde{\mu}_p,$$

with $d\tilde{\mu}_p$ the p-adic measure on $\mathcal{X}(R)$. It is enough to check that $N(\tilde{\mu}(\mathrm{ord}_\varpi(f) = n))$ is equal to the p-adic measure of the set of points x of $\mathcal{X}(R)$ with $\mathrm{ord}_\varpi(f)(x) = n$, which follows from Lemma 4.6.2. □

4.6.2. Lemma. — *Let K be a local field with residue field $k = \mathbf{F}_q$. Let $\mathcal{X}$ be a smooth formal R-scheme of dimension d. Let A be a (stable) cylinder in* $\mathrm{Gr}(\mathcal{X})$. *Then $N(\tilde{\mu}(A))$ is equal to the p-adic volume of $A \cap \mathrm{Gr}(\mathcal{X})(k)$.*

Proof. — Write $A = \pi_n^{-1}(C)$, with C a constructible subset if $\mathrm{Gr}_n(\mathcal{X})$. By definition $\tilde{\mu}(A) = \mathbf{L}^{-d(n+1)}[C]$. On the other hand $\mathcal{X}$ being smooth, the morphism $A \cap \mathrm{Gr}(\mathcal{X})(k) \to C(k)$ is surjective and its fibres are balls of radius $q^{-d(n+1)}$. It follows that the p-adic volume of $A \cap \mathrm{Gr}(\mathcal{X})(k)$ is equal to $|C(k)|q^{-d(n+1)}$. □

Let us now explain the relation with the work of Serre in **[25]**. Serre shows in **[25]** that any compact locally analytic smooth variety over K of pure dimension d is isomorphic to rB^d, with r an integer ≥ 1 and B^d the unit ball of dimension d and that, futhermore, rB^d is isomorphic to $r'B^d$ if and only if r and r' are congruent modulo $q-1$. We shall denote by $s(X)$ the class of r in $\mathbf{Z}/(q-1)\mathbf{Z}$. It follows from Serre's results that s induces an isomorphism $s : K_0(\mathrm{SLocAn}_K^d) \to \mathbf{Z}/(q-1)\mathbf{Z}$ and that the diagram

$$\begin{tikzcd}
K_0(\mathrm{GSLocAn}_K^d) \arrow[r, "\mathrm{int}_p"] \arrow[d] & \mathbf{Z}[q^{-1}] \arrow[d] \\
K_0(\mathrm{SLocAn}_K^d) \arrow[r, "s"] & \mathbf{Z}/(q-1)\mathbf{Z}
\end{tikzcd}$$

is commutative. The following result then follows from Proposition 4.6.1.

4.6.3. Corollary. — *Let K be a local field with residue field $k = \mathbf{F}_q$. Then the diagram*

$$\begin{tikzcd}
K_0(\mathrm{SRig}_K^d) \arrow[r, "\lambda"] \arrow[d, "F"] & K_0(\mathrm{Var}_k)_{\mathrm{loc}}/(\mathbf{L}-1)K_0(\mathrm{Var}_k)_{\mathrm{loc}} \arrow[d, "N"] \\
K_0(\mathrm{SLocAn}_K^d) \arrow[r, "s"] & \mathbf{Z}/(q-1)\mathbf{Z}
\end{tikzcd}$$

is commutative. □

5. Essential components of weak Néron models

5.1. Essential components and the Nash problem. — Since we shall proceed by analogy with [**21**], let us begin by recalling some material from that paper. We assume in this subsection that k is of characteristic zero and that $R = k[[\varpi]]$. For X an algebraic variety over k, we denote by $\mathcal{L}(X)$ its arc space as defined in [**10**]. In fact, in the present §, we shall use notations and results from [**10**], even when they happen to be special cases of ones in this paper. As remarked in 2.3.3, $\mathcal{L}(X) = \mathrm{Gr}(X \widehat{\otimes} R)$ and there are natural morphisms $\pi_n : \mathcal{L}(X) \to \mathcal{L}_n(X)$ with $\mathcal{L}_n(X) = \mathrm{Gr}_n(X \otimes R_n)$.

By a desingularization of a variety X we mean a proper and birational morphism

$$h : Y \longrightarrow X,$$

with Y a smooth variety, inducing an isomorphism between $h^{-1}(X \setminus X_{\mathrm{sing}})$ and $X \setminus X_{\mathrm{sing}}$ (some authors omit the last condition).

Let $h : Y \to X$ be a desingularization of X and let D be an irreducible component of $h^{-1}(X_{\mathrm{sing}})$ of codimension 1 in Y. If $h' : Y' \to X$ is another desingularization of X, the birational map $\phi : h'^{-1} \circ h : Y \dashrightarrow Y'$ is defined at the generic point ξ of D, since h' is proper, hence we can define the image of D in Y' as the closure of $\phi(\xi)$ in Y'. One says that D is an essential divisor with respect to X, if, for every desingularization $h' : Y' \to X$ of X the image of D in Y' is a divisor and that D is an essential component with respect to X, if, for every desingularization $h' : Y' \to X$ of X the image of D in Y' is an irreducible component of $h'^{-1}(X_{\mathrm{sing}})$. In general, if D is an irreducible component of $h^{-1}(X_{\mathrm{sing}})$, we say D is an essential component with respect to X, if there exists a proper birational morphism $p : Y' \to Y$, with Y' smooth, and a divisor D' in Y' such that D' is an essential component with respect to X and $p(D') = D$. It follows from the definitions and Hironaka's Theorem that essential components of different resolutions of the same variety X are in natural bijection, hence we may denote by $\tau(X)$ the number of essential components in any resolution of X.

Let W be a constructible subset of an algebraic variety Z. We say that W is irreducible in Z if the Zariski closure $\overline{W}$ of W in Z is irreducible.

In general let $\overline{W} = \cup_{1 \leq i \leq n} W'_i$ be the decomposition of $\overline{W}$ into irreducible components. Clearly $W_i := W'_i \cap W$ is non empty, irreducible in Z, and its closure in Z is equal to W'_i. We call the W_i's the irreducible components of W in Z.

Let E be a locally closed subset of $h^{-1}(X_{\mathrm{sing}})$. We denote by Z_E the set of arcs in $\mathcal{L}(Y)$ whose origin lies on E but which are not contained in E. In other words $Z_E = \pi_0^{-1}(E) \setminus \mathcal{L}(E)$. Let us remark that if E is smooth and connected, $\pi_n(Z_E)$ is constructible and irreducible in $\mathcal{L}_n(Y)$. Now we set $N_E := h(Z_E)$. Since $\pi_n(N_E)$ is the image of $\pi_n(Z_E)$ under the morphism $\mathcal{L}_n(Y) \to \mathcal{L}_n(X)$ induced by h, it follows that $\pi_n(N_E)$ is constructible and irreducible in $\mathcal{L}_n(Y)$.

The following result, proved in [**21**], follows easily from the above remarks and Hironaka's resolution of singularities:

***5.1.1. Proposition* (Nash [21]).** — *Let X be an algebraic variety over k, a field of characteristic zero. Set $\mathcal{N}(X) := \pi_0^{-1}(X_{\mathrm{sing}}) \setminus \mathcal{L}(X_{\mathrm{sing}})$. For every $n \geq 0$, $\pi_n(\mathcal{N}(X))$ is a constructible subset of $\mathcal{L}_n(X)$. Denote by W_n^1, $\ldots W_n^{r(n)}$ the irreducible components of $\pi_n(\mathcal{N}(X))$. The mapping $n \mapsto r(n)$ is nondecreasing and bounded by the number $\tau(X)$ of essential components occuring in a resolution of X.*

Up to renumbering, we may assume that W_{n+1}^i maps to $\overline{W_n^i}$ for $n \gg 0$. Let us call the family $(W_n^i)_{n\gg 0}$ a Nash family. Nash shows furthermore that for every Nash family $(W_n^i)_{n\gg 0}$ there exists a unique essential component E in a given resolution $h : Y \to X$ of X such that $\overline{\pi_n(N_E)} = \overline{W_n^i}$ for $n \gg 0$.
Now, we can formulate the Nash problem:

***5.1.2. Problem* (Nash [21] p.36).** — *Is there always a corresponding Nash family for an essential component? In general, how completely do the essential components correspond to Nash families? What is the relation between $\tau(X)$ and $\lim r(n)$?*

Let W be a constructible subset of some variety X. We denote the supremum of the dimension of the irreducible components of the closure of W in Y by $\dim W$.
Let $h : Y \to X$ be a proper birational morphism with Y a smooth variety. Let E be a codimension 1 irreducible component of the exceptional locus of h in Y. We denote by $\nu(E) - 1$ the length of $\Omega_Y^d / h^*\Omega_X^d$ at the generic point of E. Here Ω_X^d denotes the d-th exterior power of the sheaf Ω_X^1 of differentials on X.

***5.1.3. Proposition*.** — *Let X be a variety of pure dimension d over k, a field of characteristic 0. Let $h : Y \to X$ be a proper birational morphism with Y a smooth variety and let U be a non empty open subset of a codimension 1 irreducible component E of the exceptional locus of h in Y. Then*

$$\dim \pi_n(N_U) = (n+1)\, d - \nu(E)$$

for $n \gg 0$.

Proof. — By Theorem 6.1 of **[10]**, the image of $[\pi_n(N_U)]\mathbf{L}^{-(n+1)d}$ in $\widehat{K_0(\mathrm{Var}_k)}$ converges to $\mu(N_U)$ in $\widehat{K_0(\mathrm{Var}_k)}$. Since $\dim \pi_n(N_U) \leq (n+1)d$ by Lemma 4.3 of **[10]**, one deduces the fact that $\dim \pi_n(N_U) - (n+1)\, d$ has a limit. To conclude we first remark that $\overline{\pi_n(N_U)} = \overline{\pi_n(N_E)}$ for any non empty open subset U in E. Hence we may assume that $(h^*\Omega_X^d)/\mathrm{torsion}$ is locally free on a neighborhood of U. It then follows from Proposition 6.3.2 in **[10]**, or rather from its proof, that

$$\mu(N_U) = \mathbf{L}^{-d}[U](\mathbf{L}-1)\sum_{\ell\geq 1} \mathbf{L}^{-\ell\nu(E)}$$

in $\widehat{K_0(\mathrm{Var}_k)}$. Hence $\mu(N_U)$ belongs to $F^{\nu(E)}$ and not to $F^{\nu(E)+1}$, and the result follows. □

5.2. Essential components of weak Néron models. — We shall return now to the setting of the present paper. We shall fix a flat formal R-scheme $\mathcal{X}$ of relative dimension d with smooth generic fibre $\mathcal{X}_K$. By a weak Néron model of $\mathcal{X}$, we shall mean a weak Néron model $\mathcal{U}$ of $\mathcal{X}_K$ together with a morphism $h : \mathcal{U} \to \mathcal{X}$ inducing the inclusion $\mathcal{U}_K \hookrightarrow \mathcal{X}_K$. As before we shall denote by U_0^i, $i \in J$, the irreducible components of the special fibre of $\mathcal{U}$. Let ξ^i denote the generic point of U_0^i. We shall say U_0^i is an essential component with respect to $\mathcal{X}$ if, for every weak Néron model $\mathcal{U}'$ of $\mathcal{X}$, the Zariski closure of $\pi_{0,\mathcal{U}'}(\pi_{0,\mathcal{U}}^{-1}(\xi^i))$ is an irreducible component of the special fibre of $\mathcal{U}'$. Note that being an essential component is a property relative to $\mathcal{X}$. By their very definition, essential components in different weak Néron models of $\mathcal{X}$ are in natural bijection.
We have the following analogue of Proposition 5.1.1.

5.2.1. Proposition. — *Let $\mathcal{X}$ be a flat formal R-scheme of relative dimension d with smooth generic fibre $\mathcal{X}_K$. Denote by $W_n^1, \ldots W_n^{r(n)}$ the irreducible components of the constructible subset $\pi_n(\mathrm{Gr}(\mathcal{X}))$ of $\mathrm{Gr}_n(\mathcal{X})$. The mapping $n \mapsto r(n)$ is nondecreasing and bounded by the number $\tau(\mathcal{X})$ of essential components occuring in a weak Néron model of $\mathcal{X}$.*

Proof. — Clearly the mapping $n \mapsto r(n)$ is nondecreasing. Let $h : \mathcal{U} \to \mathcal{X}$ be a weak Néron model of $\mathcal{X}$, with irreducible components $\mathcal{U}^i$, $i \in J$. Since $\mathcal{U}^i$ is smooth and irreducible, $\pi_{n,\mathcal{U}}(\mathrm{Gr}(\mathcal{U}^i))$ is also smooth and irreducible, hence the Zariski closure of $h(\pi_{n,\mathcal{U}}(\mathrm{Gr}(\mathcal{U}^i))) = \pi_{n,\mathcal{X}}(\mathrm{Gr}(\mathcal{U}^i))$ in $\mathrm{Gr}_n(\mathcal{X})$ is irreducible. Since $\mathrm{Gr}(\mathcal{X})$ is the union of the subschemes $\mathrm{Gr}(\mathcal{U}^i)$, it follows that $r(n)$ is bounded by $|J|$. Now if $\mathcal{U}^i$ is not an essential component, there exists some weak Néron model of $\mathcal{X}$, $h' : \mathcal{U}' \to \mathcal{X}$, such that, if we denote by W^i the image of $\mathrm{Gr}(\mathcal{U}^i)$ in $\mathrm{Gr}(\mathcal{U}')$, $\pi_{n,\mathcal{U}'}(W^i)$ is contained in the Zariski closure of $\pi_{n,\mathcal{U}'}(\mathrm{Gr}(\mathcal{U}') \setminus W^i)$. It follows that $\pi_{n,\mathcal{X}}(\mathrm{Gr}(\mathcal{U}^i))$ is contained in the closure of $\pi_{n,\mathcal{X}}(\mathrm{Gr}(\mathcal{U}) \setminus \mathrm{Gr}(\mathcal{U}^i))$. The bound $r(n) \leq \tau(\mathcal{X})$ follows. □

As previously, we may, up to renumbering, assume that W_{n+1}^i maps to $\overline{W_n^i}$ for $n \gg 0$. We shall still call the family $(W_n^i)_{n \gg 0}$ a Nash family. Let ξ_n^i be the generic point of $\overline{W_n^i}$. By construction ξ_{n+1}^i maps to ξ_n^i under the truncation morphism $\mathrm{Gr}_{n+1}(\mathcal{X}) \to \mathrm{Gr}_n(\mathcal{X})$, hence to the inverse system $(\xi_n^i)_{n \gg 0}$ corresponds a point ξ^i of $\mathrm{Gr}(\mathcal{X})$. Let $h : \mathcal{U} \to \mathcal{X}$ be weak Néron model of $\mathcal{X}$ with irreducible components $\mathcal{U}^j$, $j \in J$. There is a unique irreducible component $\mathcal{U}^{j(i)}$ such that the point ξ^i belongs to $\mathrm{Gr}(\mathcal{U}^{j(i)})$. Furthermore, $\overline{h(\pi_n(\mathrm{Gr}(\mathcal{U}^{j(i)}))} = \overline{W_n^i}$ for $n \gg 0$ and it follows from the proof of Proposition 5.2.1 that $U_0^{j(i)}$ is essential.
We have the following analogue of the Nash problem:

5.2.2. Problem. — *Is there always a corresponding Nash family for an essential component? In general, how completely do the essential components correspond to Nash families? What is the relation between $\tau(\mathcal{X})$ and $\lim r(n)$?*

For E a locally closed subset of the special fibre of $\mathcal{U}$, we set $Z_E := \pi_{0,\mathcal{U}}^{-1}(E))$ and $N_E := h(Z_E)$. Remark that, for every n, $\pi_n(Z_E)$ and $\pi_n(N_E)$ are constructible

subsets of $\mathrm{Gr}_n(\mathcal{U})$ and $\mathrm{Gr}_n(\mathcal{X})$, respectively. Indeed, $\pi_n(Z_E)$ is constructible since $\mathcal{U}$ is smooth, hence $\pi_n(N_E) = h(\pi_n(Z_E))$ also. We denote by $\nu(U_0^i) - 1$ the length of $\Omega^d_{\mathcal{U}|R}/h^*\Omega^d_{\mathcal{X}|R}$ at the generic point ξ^i of U_0^i.
We also have the following analogue of Proposition 5.1.3.

5.2.3. Proposition. — *Let $\mathcal{X}$ be a flat formal R-scheme of relative dimension d with smooth generic fibre $\mathcal{X}_K$. Let $h : \mathcal{U} \to \mathcal{X}$ be a weak Néron model of $\mathcal{X}$ and let E be an open dense subset of an irreducible component U_0^i of the special fibre of $\mathcal{U}$. Then*

$$\dim \pi_n(N_E) = (n+1)\, d - \nu(U_0^i)$$

for $n \gg 0$.

Proof. — Fix an integer $e \geq 0$. By Lemma 3.9.2, for $n \gg e$,

$$\begin{aligned}\dim \pi_n(N_E \cap \mathrm{Gr}^{(e)}\mathcal{X}) &= \dim \pi_n(h^{-1}(N_E \cap \mathrm{Gr}^{(e)}\mathcal{X})) - \nu(U_0^i)\\ &= (n+1)d - \nu(U_0^i).\end{aligned}$$

On the other hand, it follows from Lemma 3.3.2 that

$$\dim \pi_n(N_E \cap (\mathcal{X} \setminus \mathrm{Gr}^{(e)}\mathcal{X})) < (n+1)d - \nu(U_0^i),$$

when $n \gg e \gg \nu(U_0^i)$. □

References

[1] V. Batyrev, *Birational Calabi-Yau n-folds have equal Betti numbers*, New trends in algebraic geometry (Warwick, 1996), London Math. Soc. Lecture Note Ser., 264, Cambridge Univ. Press, (1999) 1–11.

[2] V. Batyrev, *Stringy Hodge numbers of varieties with Gorenstein canonical singularities*, Proc. Taniguchi Symposium 1997, "Integrable Systems and Algebraic Geometry, Kobe/Kyoto", World Scientific, (1998) 1–32.

[3] V. Batyrev, *Non-archimedian integrals and stringy Euler numbers of log terminal pairs*, Journal of European Math. Soc. **1** (1999), 5-33.

[4] S. Bosch, U. Güntzer, R. Remmert, *Non-Archimedean analysis*, Grundlehren Math. Wiss. **261**, Springer-Verlag, Berlin (1984).

[5] S. Bosch, W. Lütkebohmert, M. Raynaud, *Néron Models*, Ergeb. Math. Grenzgeb. (3) **21**, Springer-Verlag, Berlin (1990).

[6] S. Bosch, W. Lütkebohmert, *Formal and rigid geometry. I. Rigid spaces*, Math. Ann. **295** (1993), 291–317.

[7] S. Bosch, W. Lütkebohmert, *Formal and rigid geometry. II. Flattening techniques*, Math. Ann. **296** (1993), 403–429.

[8] S. Bosch, W. Lütkebohmert, M. Raynaud, *Formal and rigid geometry. III. The relative maximum principle*, Math. Ann. **302**, (1995), 1–29.

[9] S. Bosch, K. Schlöter, *Néron models in the setting of formal and rigid geometry*, Math. Ann. **301** (1995), 339–362.

[10] J. Denef, F. Loeser, *Germs of arcs on singular algebraic varieties and motivic integration*, Invent. Math. **135** (1999), 201–232.

[11] J. Denef, F. Loeser, *Motivic integration, quotient singularities and the McKay correspondence*, to appear in Compositio Math.

[12] J. Denef, F. Loeser, *Geometry on arc spaces of algebraic varieties*, Proceedings of the Third European Congress of Mathematics, Barcelona 2000 (to appear), 22 pages.

[13] J. Dieudonné, A. Grothendieck, *Eléments de Géométrie Algébrique I*, Grundlehren Math. Wiss. **166**, Springer-Verlag, Berlin 1971.

[14] M. J. Greenberg, *Schemata over local rings*, Ann. of Math. **73** (1961), 624–648.

[15] M. J. Greenberg, *Rational points in Henselian discrete valuation rings*, Inst. Hautes Études Sci. Publ. Math. **31** (1966), 59–64.

[16] B. H. Gross, *On the motive of a reductive group*, Invent. Math. **130** (1997), 287–313.

[17] M. Kontsevich, Lecture at Orsay (December 7, 1995).

[18] M. Kontsevich, Y. Soibelman, *Homological mirror symmetry and torus fibrations*, math.SG/0011041.

[19] E. Looijenga, *Motivic measures*, in Séminaire Bourbaki, exposé 874, Mars 2000.

[20] W. Lütkebohmert, *Formal-algebraic and rigid-analytic geometry*, Math. Ann. **286** (1990), 341–371.

[21] J.F. Nash Jr., *Arc structure of singularities*, Duke Math. Journ. **81** (1995), 31–38.

[22] M. Raynaud, *Géométrie analytique rigide d'après Tate, Kiehl,...*, Table Ronde d'Analyse non archimédienne (Paris, 1972), Bull. Soc. Math. France, Mém. **39–40** (1974), 319–327.

[23] N. Schappacher, *Some remarks on a theorem of M. J. Greenberg*, Proceedings of the Queen's Number Theory Conference, Queen's Papers in Pure and Appl. Math. **54**, (1980), 101–113.

[24] J. Sebag, Intégration motivique sur les schémas formels, in preparation.

[25] J.-P. Serre, *Classification des variétés analytiques p-adiques compactes*, Topology **3** (1965), 409–412.

François Loeser, École Normale Supérieure, Département de mathématiques et applications, 45 rue d'Ulm, 75230 Paris Cedex 05, France (UMR 8553 du CNRS) *E-mail :* `Francois.Loeser@ens.fr` • *Url :* `http://www.dma.ens.fr/~loeser/`

Julien Sebag, École Normale Supérieure, Département de mathématiques et applications, 45 rue d'Ulm, 75230 Paris Cedex 05, France (UMR 8553 du CNRS) • *E-mail :* `Julien.Sebag@ens.fr`